VLA OBSERVING APPLICATION

A

rcvd:

DEADLINES: 1st of Feb., June., Oct. for next configuration following review
INSTRUCTIONS: Each numbered item must have an entry or N/A
E-MAIL TO: propsoc@nrao.edu
OR MAIL TO: Director NRAO, 520 Edgemont Rd., Charlottesville, VA 22903-2475

(1) Date Prepared: October 10, 2004

(2) Title of Proposal: **An X-ray/Radio Transient 2.5″ from Sgr A***

(3) AUTHORS (Add * for new location)	INSTITUTION	E-mail	Grad Students Only: For Ph.D. Thesis?	Grad Students Only: Anticipated Ph.D. Year
Geoffrey C. Bower	UC Berkeley	gbower@astron.berkeley.edu		
Mike Muno	UCLA	mmuno@astro.ucla.edu		
Miller Goss	NRAO	mgoss@nrao.edu		
D.C. Backer	UC Berkeley	dbacker@astro.berkeley.edu		
Yoram Lithwick	yoram@astro.berkeley.edu			
Robin Herrnstein	Columbia	herrnstein@astro.columbia.edu		
H. Falcke	ASTRON	falcke@astron.nl		

(4) Related VLA previous proposal number(s): AB1134, AB1167

(5) Contact author
for scheduling: Geoffrey C. Bower
address: 601 Campbell Hall
Astronomy Dept.
University of California, Berkeley,
CA 94720.

(6) Telephone: 510 642 4075
E-mail: gbower@astron.berkeley.edu
Fax: 510 642 3411

(7) Scientific Category: ◯ solar system ⊗ galactic ◯ extragalactic ◯ other:

(8) Configurations (one per column) (A+Pt, A, B, C, D, BnA, CnB, DnC, Any)	A				
(9) Wavelength(s) (400, 90, 20, 6, 3.5, 2, 1.3, 0.7 cm)	20,6,3.6,1.3				
(10) Time requested (hours)	3 x 1				

(11) Type of observation: ⊗ continuum ◯ spectroscopy ◯ multichannel continuum ⊗ polarimetry ◯ solar
(check all that apply) ◯ pulsar ◯ high-time resolution ◯ Pie Town link ◯ other:

(12) Suitable for dynamic scheduling? ◯ Suitable ⊗ Unsuitable

(13) ABSTRACT (do not write outside this space)

We have detected a transient radio source within 2.5 arcsec of Sgr A*. VLA measurements of the spectrum and its evolution are critical for studying radio/X-ray luminosity correlations, supporting VLBA observations, and potentially detecting diffuse structure. This source has the potential for major advances in Galacter Center science related to stellar remnants, interaction of nonthermal and thermal matter, and scattering.

NRAO use only
(03/02)

(14) Observer present for observations? ◯ Yes ⊗ No Data analysis at? ⊗ Home ◯ AOC or CV (2 weeks notice)

(15) Help required: ◯ None ◯ Consultation ◯ Friend (extensive help)

(16) Spectroscopy only	line 1	line 2	line 3	line 4
Transition (HI, OH, etc.)				
Rest Frequency (MHz)				
Velocity (km/s)				
Observing frequency (MHz)				
Correlator mode				
IF bandwidth(s) (MHz)				
Hanning smoothing (y/n)				
Number of channels per IF				
Frequency Resolution (kHz/channel)				
Rms noise (mJy/bm, nat. weight., 1 hr)				
Rms noise (K, nat. weight., 1 hr)				

(17) Number of sources:	**1**

(If more than 10 please attach list. If more than 30 give only selection criteria and LST range(s).)

(18) NAME	Coordinates 1950 ◯ 2000 ⊗ RA hh mm Dec. ± xx.x°	Conf.	λ (cm)	Corr. mode	Band-width per IF (MHz)	Total Flux (Jy)*	LAS	Required rms (mJy/bm)	Required dynamic range	Time request (hr)
CXO J174540.0-290031	17 45, -29.0	A	20,6,4,1.3		50	0.030	point source	0.1	100	

*For spectral line, this should be the total flux at the peak of the line

Notes to the table (if any):

(19) Restrictions to elevation (other than hardware limits) or HA range (give reason): We request 18 hours centered at 12 hours LST.

(20) Preferred range of dates for scheduling (give reason):
Target of opportunity!

(21) Dates which are not acceptable:

(22) Special hardware, software, or operating requirements:

(23) Please attach a self-contained Scientific Justification **not in excess of 1000 words.** (Preprints or reprints will be ignored.)
Please include the full addresses (postal and e-mail) for first-time users or for those that have moved (if not contact author).
When your proposal is scheduled, the contents of the cover sheets become public information (Any supporting pages are for refereeing only).

An X-ray/Radio Transient 2.5″ from Sgr A*

Bower et al.

1. Transient Discovery

In observations on 1 and 4 October 2004, we detected a radio transient in the Galactic Center (GC) at L-band with the VLA. Observations at 1515 MHz and 1715 MHz on 1 October 2004 and 1365 MHz on 4 October 2004 show a ~ 30 mJy source 2.5″ south of Sgr A* (Figure 1). This source is not present in archival (24 June 1999) L-band observations of Sgr A* (Figure 2).

Follow-up observations on 7 October 2004 show that the source continues to be present at L band as well as appearing at 8.4 GHz and 22 GHz (Figure 3). We measure flux densities of 30 ± 5 mJy, 10 ± 2 mJy, and 4 ± 1 mJy at 1.4, 8.4 and 22 GHz, respectively. Imaging of this source is particularly difficult because it lies very close to the bright point source Sgr A* and it lies within a region dominated by diffuse flux. We achieve a position with an accuracy of 0.02 arcsec with the 8.4 GHz observations.

The radio source is positionally coincident with an X-ray transient (CXO J174540.0-290031; Figure 4). The best fit radio position is 17:45:40.063, -29:00:30.622, in agreement at < 0.4 arcsec with the X-ray position, which has an accuracy of ~ 0.5 arcsec. The X-ray transient was bright during XMM observations in April, and Chandra observations in July and at the end of August. It brightened by a factor of 2 between July-September. The source started out hard in July, with a spectrum that could be described as a power-law with photon index $\Gamma = 0$. By September, the source had softened to a $\Gamma = 1.0$ power law, which is still hard, but more typical of an accreting neutron star or black hole. Its luminosity of $> 10^{34}$ erg/s indicates that it is an accreting black hole or neutron star.

The X-ray luminosity is quite low, $\sim 10^{-5}$ to $10^{-4} L_{Edd}$, however, for a compact object with strong radio emission. The ratio of peak X-ray to peak radio emission is $\sim 2 \times 10^5$ mJy/crab, which is rivaled only by Cyg X-3 (Fender & Kuulkers 2001) with a ratio of 2×10^4. Typical values of this ratio are in the range 100 to 1000. RXTE ASM observations would have caught a recent X-ray burst as faint as $10^{-2} L_{Edd}$.

The precise VLA position for the radio source permits us to determine that there is no infrared counterpart at a limit of $K = 16.5$ (Lu & Ghez, private communication). This indicates that the companion of the transient source is fainter than a B-star.

The diffuse X-ray emission within 3" of the transient has brightened by a factor of 2

over previous observations. This may be reflected X-ray emission, but it could also be energy input from the action of a jet.

This source offers an exciting and rare opportunity to study a compact object in the GC. This one of only a handful of GC transient radio sources and it is by far the closest to Sgr A* (e.g., Davies et al. 1976; Zhao et al. 1986; Hyman et al. 2002). We know very little about the compact object population of the GC, its relationship to massive star formation, and bulge formation (Miralde-Escude & Gould 2000; Hansen & Milosavljevic 2003). Potential science includes:

- A study of the radio/X-ray luminosity. The quite large radio to X-ray luminosity ratio indicates that this object may be quite unususal. A better characterization of the radio spectrum and its evolution will help us to compare this source to other galactic X-ray sources (e.g., Fender, Belloni & Gallo 2004) and extragalactic sources (Falcke, Körding & Markoff 2004).

- A high resolution radio image of the X-ray transient source. Detection of a jet may resolve the question of the nature of this object and the diffuse X-ray emission.

- Proper motion with respect to Sgr A*. A small PM with respect to Sgr A* will confirm that the source is galactic; an extragalactic background source would show a PM of 6 mas/y. Additionally, we can ask: Is it in a bound orbit? Does it share a similar velocity with the high mass stars in the neighborhood? Typical stellar velocity dispersion at this radius from Sgr A* is 150 ± 50 km/s (Ghez et al. 1998) or 10 microarcsec/day. Compact objects such as pulsars often have proper motions of ~ 300 km/s, much higher than field stars (Brisken et al. 2003). If this object persists for a month or more, we can measure its proper motion with multiple VLBA observations. The transient studied by Zhao et al. (1986) persisted for ~ 6 months before fading away.

- A new probe of scattering very close to Sgr A*. Does it have the same scattering properties as Sgr A*? This is the closest compact radio source to Sgr A*. The Zhao et al. transient at a distance of ~ 36 arcsec showed a similarity with Sgr A* in scattering properties, although the inclination of the scattering disk and its axial ratio were not determined. Variations in these parameters has implications for the magnetic field structure and the distribution of ionized gas in the GC.

2. Requested Observations

2.1. VLA

Continued radio monitoring of this object is essential for determining the suitability of VLBA measurements as well as for characterizing the radio/X-ray luminosity correlation. We are pursuing further X-ray observations. We request three observations over the next two weeks. We will measure the spectrum at 1.4, 5.0, 8.4 and 22 GHz. We require 60 minutes per epoch to give us a range in (u,v) coverage at each frequency that will permit us to resolve out the diffuse structure.

2.2. VLBA

We request two VLBA observations, spaced by ~ 10 days, as soon as possible. A proper motion a few times in excess of the stellar velocity dispersion will be detectable over this span.

While the L-band flux density appears to be stable over the past 10 days, the flux density at higher frequencies may fall off more rapidly with time. The steep spectral index of this source suggests that it may be fading. Observations will be performed at 8.4 GHz and 22 GHz. The two frequencies will allow us to characterize the angular size as a function of wavelength, which is important for scattering studies and for determining the intrinsic structure.

Because of the extreme scattering environment, we will not need HN, SC or MK. Baselines to these stations are resolved out. With 1 hour of observation at 8.4 GHz and 3 hours of observation at 22 GHz, we can achieve an image rms of 200 μJy and 320 μJy and resolution (set by Sgr A* scattering size) at 17 mas and 2.4 mas at the two frequencies, respectively. This will enable us to achieve $> 10\sigma$ detections and source positions to an accuracy of ~ 1 mas and ~ 0.1 mas, respectively.

We have verified the methods of this observation through use of a March 2004 VLBA 43 GHz observation of Sgr A* (BB183). We attain an rms of 800 μJy at the location of the transient, which is close to the expected noise for these observations.

Fig. 1.— *(left)* 1715 MHz image obtained on 1 October 2004. Sgr A* is the bright source and the transient is due south of it. Beam size is 1.5 x 0.5 arcsec in p.a. 10 deg. *(right)* 1365 MHz image obtained on 4 October 2004. Images were made with inner (u,v) cutoff of 50 kiloλ. Beam size is 2.1 x 0.8 arcsec in p.a. -3 deg. 1715 MHz image included PT data.



3. References

Brisken, W., et al., 2003, AJ, 126, 3090
Davies, R.D., et al., 1976, Nature, 261, 476
Falcke, H., Körding, E., & Markoff, S., 2004, A&A, 414, 895
Fender, R.P. & Kuulkers, E., 2001, MNRAS, 324, 923
Fender,R.P., Belloni, T.M., & Gallo, E., 2004, A&A, submitted, astro-ph/0409360
Ghez, A.M., et al., 1998, ApJ, 509, 678
Hansen, B.M.S. & Milosavljevic, M., 2003, ApJL, 593, L77
Hyman, S.D., et al., 2002, AJ, 123, 1497
Miralde-Escude, J. & Gould, A., 2000, ApJ, 545, 847
Zhao, J.-H., et al., 1992, Science, 255, 1538

Fig. 2.— 1365 MHz observation of Sgr A* from 24 June 1999, showing no transient source. Beam size is 2.3 x 0.9 arcsec in p.a. 6 deg.



Fig. 3.— 8.4 GHz observation of Sgr A*, showing the transient source. Beam size is 0.44 x 0.16 arcsec in p.a. 12 deg.



Fig. 4.— CXO X-ray image before and after appearance of the transient.